

Tying in High Resolution E3SM with ARM Data (THREAD)

Lawrence Livermore National Laboratory

ARM





Storm Resolving

Regionally Refined SCREAM (RRM-SCREAM) for ARM Sites

August 9, 2023

Hsi-Yen Ma

and

Yunyan Zhang, Peter Bogenschutz, Jishi Zhang Lawrence Livermore National Laboratory

> This work was performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under contract DF-AC52-07NA27344_LLNL-PRES-838692



RRM-SCREAM

- RRM: An effective and efficient tool for high-resolution model development and diagnosis
- Six RRM-SCREAM configurations are proposed to study convection over land, marine low clouds and landatmosphere interactions.
- Inner domain (refined region) will be ~3.25 km or 1.6 km, and will be freerunning. Outer domain will be ~100 km, and the dynamical fields in the outer domain will be nudged toward the ERA5 reanalysis

RRM-SCREAM configurations



Validation of RRM-SCREAM simulations









Experiments:

- SCREAMv0, starting from 2020/01/20, 40-day long (DYAMOND II)
- 2 RRM-SCREAM, 5-day long hindcasts starting from 2020/01/20 (w/ and w/o nudging)

Transition of shallow to deep convection (GoAmazon)





Case study:

- Transition of shallow to deep convection:
 - Single-peak days: 2014/6/27, 2014/10/05
 - Double-peak days: 2014/8/11, 2015/8/26





- Relevant to our current GoAmazon model simulations and analysis:
 - —How can we better utilize ARM data to compare with km-scale model output (making meaningful comparison, e.g., using instrument simulators)?
 - -What are the limitation in applying these data?
- We are looking for ideas on how to better utilize RRM-SCREAM configuration for relevant cloud, convection and L-A interactions with ARM data:
 —What other field campaign locations would you mostly be interested in using RRM-SCREAM for modeling studies (e.g., AMF3 BNF, EPCAPE)?